

Generalized Quantum Field Theory as an Alternative Approach To The Problem of Composite Particles Reaction

C.I. Ribeiro-Silva and N. M. Oliveira-Neto
 Centro Brasileiro de Pesquisas Físicas,
 Rua Xavier Sigaud, 150 cep 22290-180,
 Rio de Janeiro, Brazil

A generalization of the Heisenberg algebra has been recently constructed. This generalized algebra has a characteristic function which depends on one of its generators. When this function is linear, $qJ_0 + s$, it is possible to construct a Generalized Quantum Field Theory (GQFT) that creates at a space-time a composite particle. In the present work we show that a generalized QFT can also be constructed consistently, even with a nonlinear characteristic function and leads to better results as long as we have more parameters to (possibly) fit the spectrum of a real composite particle.

PACS numbers: 05.40.-a, 05.90.+y, 89.65.Gh

I. INTRODUCTION

It is well known that the standard quantum field theory (QFT) is constructed within the framework of Heisenberg algebra [1,2]. Therefore a possible way to construct a non-standard QFT is through generalization of the Heisenberg algebra. A generalized Heisenberg algebra (GHA) has recently been proposed [3]. This generalized algebra has a characteristic function which depends on one of its generators. When this function is linear with slope q , the algebra turns into a q -oscillator [3] and it was shown [4] that, for this case, it is possible to construct a generalized QFT that describes at a space-time a composite particle. In Ref [4] the propagator, the first and second order scattering processes were computed and it was shown that the convergence of the perturbative series can be changed. However, other realizations of the GHA have been presented in [3,5], the case where we have a nonlinear characteristic function. We will show, in this present work, that a self-consistent GQFT can also be constructed in that case and as a consequence we have more parameters to fit experimental data. A detailed discussion comparing these two case will be presented ahead.

The first step is to analyze the GHA for a general characteristic function $f(J_0)$. As defined previously [3], this generalized Heisenberg algebra is generated by three operators, J_0, A and A^\dagger , and described by the following relations

$$J_0 A^\dagger = A^\dagger f(J_0) \quad (1)$$

$$A J_0 = f(J_0) A \quad (2)$$

$$[A, A^\dagger] = f(J_0) - J_0 \quad (3)$$

where † means Hermitian conjugate and by hypothesis, $J_0^\dagger = J_0$ and $f(J_0)$ is an arbitrary function of J_0 . Within this algebra we verify that the generators satisfy the Jacobi identity with

$$C = A^\dagger A - J_0 = A A^\dagger - f(J_0) \quad (4)$$

being the Casimir operator of the algebra. Assuming that there is a vacuum state represented by $|0\rangle$, it can be demonstrated [3] that for an arbitrary function f , that

$$J_0 |m-1\rangle = f^{(m-1)}(\alpha_0) |m-1\rangle, \quad m = 1, 2 \quad (5)$$

$$A^\dagger |m-1\rangle = N_{m-1} |m\rangle \quad (6)$$

$$A |m\rangle = N_{m-1} |m-1\rangle, \quad (7)$$

where $N_{m-1}^2 = f^m(\alpha_0) - \alpha_0$, α_0 is the lowest J_0 eigenvalue and $f^m(\alpha_0)$ is the m th iteration through function $f(\alpha_0)$. This GHA describes a class of one-dimensional quantum systems characterized by energy eigenvalues given by

$$\epsilon_n = f(\epsilon_{n-1}) \quad (8)$$

where ϵ_n and ϵ_{n-1} are successive energy levels. Unlike standard Heisenberg Algebra (where the energy of the n -th level is equal to n times the energy of the first one), here, the energy of the n -th level (depending on the value of the parameters) is greater or smaller than n times the energy of the first one. In the last case, the energy gap between consecutive levels becomes smaller as n increases, behaving like the energy spectrum of a composite particle. So we can postulate that this algebra describes a composite particle.

II. A GENERALIZED QFT

Let us discuss, firstly, the algebra given by (1)-(3) for the quadratic case, i.e. $f(J_0) = t J_0^2 + q J_0 + s$, which is the simplest nonlinear case. The algebraic relations (1)-(3) can be written as [3]

$$[J_0, A^\dagger]_q = t A^\dagger J_0^2 + s A^\dagger, \quad (9)$$

$$[J_0, A]_{q^{-1}} = -\frac{t}{q} J_0^2 A - \frac{s}{q} A, \quad (10)$$

$$[A^\dagger, A] = t J_0^2 + (q-1) J_0 + s, \quad (11)$$

where $[a, b]_q = ab - qba$, is the q -deformed commutation relation of two operators a and b . The relations (9)-(11) describe a two-parameter deformed Heisenberg Algebra already studied [3]. Of course, for $t = 0$ we recover the linear case and if additionally $q = 1$, the standard Heisenberg algebra.

We focus now on the graphical analysis of the function $f(\alpha_0) = t\alpha_0^2 + q\alpha_0 + s$. Let us plot $y = f(\alpha_0)$ together with $y = \alpha_0$. In the points where lines intersect, we have $\alpha_0 = y = f(\alpha_0)$. So the intersections are precisely the fixed points.




FIG. 1. (a) $\Delta > 0$ (b) $\Delta = 0$ and (c) $\Delta < 0$, where $\Delta = (q-1)^2 + 4ts$.

Assuming $t < 0$, there are three cases to be analyzed: (a) $\Delta < 0$ (b) $\Delta = 0$ and (c) $\Delta > 0$, for $\Delta = (q-1)^2 + 4ts$ (see Fig 1). However, as depicted in (Fig. 1), depending on t, q, s and α_0 , we have different spectra. Nevertheless, we are interested in a special spectrum which can be associated to a composite particle, thus, only the case (c) for $\alpha_{min} < \alpha_0 < \alpha_{max}$ is relevant. Fig 2 shows the comparison between the quadratic and the linear cases.

As one may notice, in the linear case, there is a simple relation among energy level gaps (as n increases, the energy gap between successive levels always decreases) which make the linear case unsuitable to fit some realistic spectra, see for instance [11]. This makes the quadratic more suitable to fit the spectra of the mainstream composite particle.




FIG. 2. Comparison between the quadratic and linear cases

Now, we will realize the operators A^\dagger , A and J_0 in terms of physical operators, as in the case of the one-dimensional harmonic oscillator¹. In order to do so we will employ the formalism of non-commutative differential and integral calculus [6] which considers a one-dimensional lattice in a momentum space where the momenta are allowed to take only discrete values $p_0, p_0 + a, p_0 + 2a$, and so on, with $a > 0$. Let us introduce the momentum shift operator

$$T = 1 + a \partial_p \quad (12)$$

$$\bar{T} = 1 - a \bar{\partial}_p \quad (13)$$

where ∂_p and $\bar{\partial}_p$ are the left and right discrete derivatives that shift the momentum value by a , i.e.

$$T f(p) = f(p + a), \quad (14)$$

$$\bar{T} f(p) = f(p - a) \quad (15)$$

and satisfy

$$T \bar{T} = \bar{T} T = \hat{1}. \quad (16)$$

Introducing the momentum operator P

$$P f(p) = p f(p), \quad (17)$$

hence

$$T P = (P + a) T \quad (18)$$

$$\bar{T} P = (P - a) \bar{T}. \quad (19)$$

Now, we return to the realization, observing that, in this case we have not an explicit formula for $f^m(\alpha_0)$ as

¹where A^\dagger and A can be written as a function of p and q .

in the linear one [3], but we can still associate this two-parameter deformed Heisenberg algebra (9)-(11) to the one-dimensional lattice we have just presented. By defining an operator N such that

$$N|m\rangle = m|m\rangle, \quad (20)$$

we can write an operator $f(N, \alpha_0)$ that verifies

$$f(N, \alpha_0)|m\rangle = f^m(\alpha_0)|m\rangle, \quad (21)$$

So, using (5)-(7) we can write J_0 as

$$J_0 \equiv f(N, \alpha_0) = \alpha_N \equiv f(P/a, \alpha_0) \quad (22)$$

where we have defined $N = P/a$, with P given by (17). This implies that

$$P|m\rangle = m a|m\rangle, \quad m = 0, 1, 2, \dots \quad (23)$$

Moreover, it is easy to see [4] that

$$\bar{T}|m\rangle = |m+1\rangle, \quad m = 0, 1, 2, \dots \quad (24)$$

with \bar{T} and $T = \bar{T}^\dagger$ defined in Eqs.(14)-(15).

Now, using Eqs.(5)-(7), we finally define

$$A^\dagger = S(P) \bar{T}, \quad (25)$$

$$A = T S(P), \quad (26)$$

where

$$S(P)^2 = J_0 - \alpha_0, \quad (27)$$

being α_0 the lowest J_0 eigenvalue. Following similar steps to those used to construct a standard spin-0 QFT [7], let us define two operators

$$\chi \equiv \imath(S(P)(1 - a\bar{\partial}_p) - (1 + a\partial_p)S(P)) = -\imath(A - A^\dagger) \quad (28)$$

$$Q \equiv S(P)(1 - a\bar{\partial}_p) + (1 + a\partial_p)S(P) = A + A^\dagger, \quad (29)$$

satisfying

$$[\chi, P] = -\imath a Q \quad (30)$$

$$[P, Q] = \imath a \chi \quad (31)$$

$$[\chi, Q] = 2\imath(S^2(P) - S^2(P + a)). \quad (32)$$

At this point, we introduce an independent copy of the one-dimensional momentum lattice, we have just defined, at each point of a \vec{K} -lattice through the substitution $P \rightarrow P_{\vec{k}}$ so,

$$A_{\vec{k}}^\dagger = S_{\vec{k}} \bar{T}_{\vec{k}}, \quad (33)$$

$$A_{\vec{k}} = T_{\vec{k}} S_{\vec{k}}, \quad (34)$$

$$J_0(\vec{k}) = f^{\frac{P_{\vec{k}}}{a}}(\alpha_0). \quad (35)$$

Then, we define three field operators

$$\phi(\vec{r}, t) = \sum_{\vec{k}} \frac{1}{\sqrt{2\Omega w(\vec{k})}} (A_{\vec{k}}^\dagger e^{-\imath \vec{k} \cdot \vec{r}} + A_{\vec{k}} e^{\imath \vec{k} \cdot \vec{r}}), \quad (36)$$

$$\Pi(\vec{r}, t) = \sum_{\vec{k}} \frac{\imath w(\vec{k})}{\sqrt{2\Omega w(\vec{k})}} (A_{\vec{k}}^\dagger e^{-\imath \vec{k} \cdot \vec{r}} - A_{\vec{k}} e^{\imath \vec{k} \cdot \vec{r}}), \quad (37)$$

$$\wp(\vec{r}, t) = \sum_{\vec{k}} \sqrt{\frac{w(\vec{k})}{2\Omega}} S_{\vec{k}} e^{-\imath \vec{k} \cdot \vec{r}}, \quad (38)$$

where $w(\vec{k}) = \sqrt{\vec{k}^2 + m^2}$ and m is a real number. Using (36)-(38) we can show that the Hamiltonian

$$H = \frac{1}{2} \int d^3 \vec{r} (\Pi^2(\vec{r}, t) + u |\wp(\vec{r}, t)|^2 + \phi(\vec{r}, t) (-\vec{\nabla}^2 + m^2) \phi(\vec{r}, t)) \quad (39)$$

can be written as

$$H = \frac{1}{2} \sum_{\vec{k}} w(\vec{k}) \left[S_{\vec{k}}^2(N+1) + (1+u) S_{\vec{k}}^2(N) \right] \quad (40)$$

where $N_0^2 = t\alpha_0^2 + (q-1)\alpha_0 + 1$ and $S^2(N)$ is given by Eq.(27). Note that in the limit $t \rightarrow 0$, we recover the linear case and for $t \rightarrow 0$, $q \rightarrow 1$ ($u \rightarrow 0$), the Hamiltonian is proportional to the number operator.

The time evolution of the fields can be studied by solving Heisenberg's equation for $A_{\vec{k}}^\dagger, A_{\vec{k}}$. So, using Eq. (40) we have

$$[H, A_{\vec{k}}^\dagger] = w(\vec{k}) A_{\vec{k}}^\dagger h(N_{\vec{k}}) \quad (41)$$

where for the quadratic case

$$h(N_{\vec{k}}) = \frac{1}{2} \Delta E \left[t(S_{\vec{k}}^2(N+1) + S_{\vec{k}}^2(N)) + 2t\alpha_0 + Q \right], \quad (42)$$

with $Q = 1 + u + q$ and $\Delta E = S_{\vec{k}}^2(N+1) - S_{\vec{k}}^2(N)$. For the general case $f(\alpha_0) = \sum_{j=0}^n a_j \alpha_0^j$, we have

$$h(N_{\vec{k}}) = \frac{1}{2} \Delta E \sum_{j=1}^n \sum_{i=0}^{j-1} \left[a_j (S_{\vec{k}}^2(N) + \alpha_0)^{j-1} \chi(N_{\vec{k}})^i \right] + (u+1) \quad (43)$$

with

$$\chi(N) = \frac{S_{\vec{k}}^2(N+1) + \alpha_0}{S_{\vec{k}}^2(N) + \alpha_0}. \quad (44)$$

Solving the Heisenberg equation

$$A_{\vec{k}}^\dagger = A_{\vec{k}}^\dagger(0) e^{\imath w(\vec{k}) h(N_{\vec{k}}) t} \quad (45)$$

we can write Eq.(36) as $\phi(\vec{r}, t) = \alpha(\vec{r}, t) + \alpha^\dagger(\vec{r}, t)$,² where

$$\alpha^\dagger(\vec{r}, t) = \sum_{\vec{k}} \frac{1}{\sqrt{2\Omega w(\vec{k})}} A_k^\dagger e^{-i\vec{k}\cdot\vec{r} + i w(\vec{k}) h(N_{\vec{k}}) t}. \quad (46)$$

The Feynman propagator $D_F^N(x_1, x_2)$ defined as Dyson-Wick contraction between $(x_i \equiv (r_i, t_i))$ $\phi(x_1)$ and $\phi(x_2)$ can be computed using (46). In the integral representation it is given by

$$D_F^N(x) = \frac{-i}{(2\pi)^4} \int d^4 k \frac{S_k^2(N+1) e^{i\vec{k}\cdot\vec{r} - i k_0 h(N_{\vec{k}}) t}}{k^2 + m^2} + (N \rightarrow N-1). \quad (47)$$

Note that, if $q \rightarrow 1$ the standard result is recovered.

III. PERTURBATIVE COMPUTATION

We shall now analyze the scattering process $1 + 2 \rightarrow 1' + 2'$ for $\vec{p}_1 \neq \vec{p}_2 \neq \vec{p}'_1 \neq \vec{p}'_2$ with initial state

$$N_0^2 |1, 2\rangle \equiv A_{\vec{p}_1}^\dagger A_{\vec{p}_2}^\dagger |0\rangle \quad (48)$$

and final state

$$N_0^2 |1, 2\rangle \equiv A_{\vec{p}_1'}^\dagger A_{\vec{p}_2'}^\dagger |0\rangle \quad (49)$$

where these particles are described by the Hamiltonian given in Eq.(40) with an interaction $\lambda \int : \phi^4(\vec{r}, t) : d^3 \vec{r}$.

For the first order scattering process we have

$$S_{fi(1)} = \frac{-6(2\pi)^4 i N_0^2 \lambda \delta^4(P_1 + P_2 - P'_1 - P'_2)}{\Omega^2 (t(N_0^2 + 2\alpha_0) + Q) \sqrt{\omega_{\vec{p}_1} \omega_{\vec{p}_2} \omega_{\vec{p}'_1} \omega_{\vec{p}'_2}}} \quad (50)$$

and for the second order

$$S_{fi(2)}^a = \frac{N_0^4 \lambda^2 \delta^4(P_1 + P_2 - P'_1 - P'_2) I}{2\Omega^2 (t(N_0^2 + 2\alpha_0) + Q)^2 \sqrt{\omega_{\vec{p}_1} \omega_{\vec{p}_2} \omega_{\vec{p}'_1} \omega_{\vec{p}'_2}}} \quad (51)$$

$$S_{fi(2)}^b = \frac{N_0^4 \lambda^2 \delta^4(P_1 + P_2 - P'_1 - P'_2) I'}{2\Omega^2 (t(N_0^2 + 2\alpha_0) + Q)^2 \sqrt{\omega_{\vec{p}_1} \omega_{\vec{p}_2} \omega_{\vec{p}'_1} \omega_{\vec{p}'_2}}} \quad (52)$$

$$S_{fi(2)}^{c,t} = \frac{N_0^4 \lambda^2 \delta^4(P_1 + P_2 - P'_1 - P'_2) I''}{8\Omega^2 (t(N_0^2 + 2\alpha_0) + Q)^2 \sqrt{\omega_{\vec{p}_1} \omega_{\vec{p}_2} \omega_{\vec{p}'_1} \omega_{\vec{p}'_2}}} \quad (53)$$

$$S_{fi(2)}^{c,u} = \frac{N_0^4 \lambda^2 \delta^4(P_1 + P_2 - P'_1 - P'_2) I'''}{8\Omega^2 (t(N_0^2 + 2\alpha_0) + Q)^2 \sqrt{\omega_{\vec{p}_1} \omega_{\vec{p}_2} \omega_{\vec{p}'_1} \omega_{\vec{p}'_2}}} \quad (54)$$

where,

²hereafter $A_k^\dagger(t=0) \equiv A_k^\dagger$

$$P_i = (\vec{p}_i, \omega_{\vec{p}_i}), \quad (55)$$

$$P'_i = (\vec{p}'_i, \omega_{\vec{p}'_i}), \quad (56)$$

$$I = \int d^4 k \frac{1}{(k^2 + m^2)((-k + s)^2 + m^2)}, \quad (57)$$

$$(58)$$

with $s = P_1 + P_2$, and

$$I' = I(s \rightarrow -s), \quad (59)$$

$$I'' = I(s \rightarrow t), \quad (60)$$

$$I''' = I(s \rightarrow u), \quad (61)$$

being

$$t = P_1 - P'_1, \quad (62)$$

$$u = P_1 - P'_2. \quad (63)$$

So, up to second order we have

$$S_{fi} = \frac{\lambda N_0^2}{(N_0^2 + 2\alpha_0) + Q} A_1 + \frac{\lambda^2 N_0^4}{((N_0^2 + 2\alpha_0) + Q)^2} (A_2^s + A_2^t + A_2^u) \quad (64)$$

where A_1, A_2^s, A_2^t and A_2^u are the same contributions that one can find in the structureless particle standard $\lambda\phi^4$ theory model corresponding to the s, t and u channels for or




FIG. 3. Comparison between the deformed cross sections for quadratic (symbols) and linear (lines) cases with the standard one.

Fig. 3 compares the deformed cross section, for the linear and quadratic cases, with the standard one [9]. As one can see, the linear case always gives a deformed-cross section smaller than the standard one for $0 < \alpha_0 < \epsilon^*$ and greater for $\alpha_0 < 0$ (for $0 < q < 1$). Therefore, it becomes difficult to fit both energy spectrum of the composite particle and cross section, inasmuch as we may have an ambiguous situation where we need $\alpha_0 < 0$ to fit the energy spectrum and $\alpha_0 > 0$ for the cross section. However, in the quadratic case we do not have such ambiguity (as depicted in Fig. 3) because we have one more parameter.

IV. CONCLUSION

We showed that within the framework of deformed Heisenberg algebra, with a quadratic characteristic function, it is possible to construct a generalized quantum field theory (GQFT) that describes a composite particle. Comparison between a GQFT made with a linear [4] and quadratic characteristic function was performed showing that, the latter, is more suitable to fit experimental data. It is worthwhile to mention that a GQFT made with a general characteristic function brings wider possibilities than the quadratic one but several restrictions must be imposed to the parameters, a_1, a_2, \dots, a_N , in order to describe a composite particle. These restrictions are already present in the quadratic case but the algebra is much more simple for this case. One can show that the general case can be obtained replacing $Q \rightarrow \sum_{j=1}^n \sum_{i=0}^{j-1} \left[a_j (\alpha_0)^{j-1} \left(\frac{N_0^2 + \alpha_0}{\alpha_0} \right)^i \right] + (u + 1)$. In future works we will address this formalism to analyze Compton scattering by nuclei below pion threshold where

no quantitative consistent description exists based on first principles [10].

Acknowledgments: The authors thank CNPq (Brazil) for partial support, Evaldo Curado and Marco Aurelio Rego-Monteiro for valuable comments

-
- [1] W. Heisenberg, Z. Physik **120**, 513, 673 (1943); Z. Naturforsch. **1**, 608 (1946).
 - [2] C. Møller, Kgl. Danske Videnskab. Selskab, Mat-fys. Medd. **23**, No 1 (1945); **24**, 19 (1946).
 - [3] E. M. F. Curado and M. A. Rego-Monteiro, J. Phys. **A 34** (2001) 3253.
 - [4] V. B. Bezerra, E. M. F. Curado and M. A. Rego-Monteiro, Phys. Rev. **D 66**, (2002), 085013.
 - [5] E. M. F. Curado, M. A. Rego-Monteiro and H. N. Nazareno, Phys. Rev. **A 64** (2001) 12105.
 - [6] A. Dimakis and F. Muller-Hoissen, Phys. Let. **B 295** (1992) 242.
 - [7] See for instance: T. D. Lee, "Particle Physics and Introduction to Field Theory", Harwood academic publishers, New York, 1981.
 - [8] D. Bonatsos, J. Phys. **A 25**, L101 (1992).
 - [9] Greiner Reinhardt, "Field Quantization", Springer-Verlag Berlin Heidelberg New York, 1996.
 - [10] M.-Th Hutt et al. **Physics Reports 323 (2000) 457-594**
 - [11] Williams, W. S. C., Nuclear and Particle Physics, Oxford University Press, 1992, pag. 67